

ADAPTATION MECHANISMS OF THE CARDIORESPIRATORY SYSTEM TO PHYSICAL LOAD IN ATHLETES: A REVIEW

Sirojiddin Omonturdiyev^{*1,3}, Alikhon Khasanov², Pulatov Laziz¹

¹Oriental University, Tashkent, Uzbekistan.

²Namangan State University, Namangan Region, Uzbekistan.

³A.S. Sadykov Institute of Bioorganic Chemistry of the Academy of Sciences of Uzbekistan, Tashkent, Uzbekistan.

Article Received: 16 March 2026 | Article Revised: 7 April 2026 | Article Accepted: 27 April 2026

*Corresponding Author: Sirojiddin Omonturdiyev

¹Oriental University, Tashkent, Uzbekistan. ²A.S. Sadykov Institute of Bioorganic Chemistry of the Academy of Sciences of Uzbekistan, Tashkent, Uzbekistan. Email ID: siroj.2012@mail.ru DOI: <https://doi.org/10.5281/zenodo.19910947>

How to cite this Article: Sirojiddin Omonturdiyev, Alikhon Khasanov, Pulatov Laziz (2026) ADAPTATION MECHANISMS OF THE CARDIORESPIRATORY SYSTEM TO PHYSICAL LOAD IN ATHLETES: A REVIEW. World Journal of Pharmaceutical Science and Research, 5(5), 63-73.



Copyright © 2026 Sirojiddin Omonturdiyev | World Journal of Pharmaceutical Science and Research.

This work is licensed under creative Commons Attribution-NonCommercial 4.0 International license (CC BY-NC 4.0).

ABSTRACT

The cardiorespiratory system plays a central role in maintaining physical performance and ensuring oxygen delivery during exercise. In athletes, repeated exposure to physical load leads to significant structural and functional adaptations in both the cardiovascular and respiratory systems. This review aims to analyze and summarize the main mechanisms of cardiorespiratory adaptation to physical exercise based on recent scientific literature. Regular physical training induces physiological changes such as increased stroke volume, reduced resting heart rate (physiological bradycardia), and enhanced cardiac output efficiency. At the respiratory level, improvements in lung ventilation, oxygen diffusion capacity, and maximal oxygen uptake (VO_{2max}) are observed. These adaptations are closely linked to enhanced capillary density, mitochondrial biogenesis, and improved oxygen utilization in skeletal muscles. Furthermore, neurohumoral regulation plays an important role in adaptation, where sympathetic-parasympathetic balance and hormonal responses (adrenaline, cortisol) modulate cardiovascular and respiratory performance during stress. The degree of adaptation varies depending on the type, intensity, and duration of physical training. Understanding these mechanisms is essential for optimizing athletic performance, designing training programs, and preventing overtraining-related disorders. This review highlights the integrated nature of cardiorespiratory adaptation and emphasizes its significance in sports physiology and medical science.

KEYWORDS: Cardiorespiratory system, physical exercise, VO_{2max} , physiological adaptation, sports physiology.

INTRODUCTION

Physical exercise represents one of the most important physiological stressors for the human body, inducing a wide range of adaptive responses across multiple organ systems. Among these, the cardiorespiratory system plays a central

role in maintaining homeostasis during increased metabolic demand. The coordinated function of the cardiovascular and respiratory systems ensures efficient oxygen delivery to working muscles and removal of metabolic by-products such as carbon dioxide. In athletes, repeated exposure to systematic physical training leads to both acute functional responses and long-term structural adaptations, which significantly enhance physical performance and endurance capacity.^[1,2]

The cardiorespiratory system includes the heart, blood vessels, lungs, and associated regulatory mechanisms that collectively support gas exchange and circulation. During physical activity, oxygen consumption increases dramatically, requiring rapid adjustments in heart rate, stroke volume, ventilation rate, and tissue perfusion. These acute responses are regulated by neural and hormonal mechanisms, primarily involving the autonomic nervous system and circulating catecholamines. However, with long-term training, the body undergoes chronic adaptations that improve efficiency at rest and during exercise.^[3,4]

One of the most significant adaptations in trained athletes is an increase in cardiac efficiency, characterized by enhanced stroke volume and reduced resting heart rate, commonly referred to as physiological bradycardia. Additionally, myocardial contractility improves, and the heart may undergo mild physiological hypertrophy, often described as “athlete’s heart.” These changes allow the cardiovascular system to deliver a greater volume of oxygenated blood with reduced energy expenditure.^[5,6]

On the respiratory side, adaptations include increased tidal volume, improved pulmonary ventilation efficiency, and enhanced diffusion capacity of the alveolar membrane. These changes contribute to a higher maximal oxygen uptake (VO_{2max}), which is widely recognized as a key indicator of aerobic fitness and endurance performance. At the muscular level, increased capillary density, mitochondrial biogenesis, and improved oxidative enzyme activity further support efficient oxygen utilization.

The importance of studying cardiorespiratory adaptations lies not only in understanding athletic performance but also in its broader implications for health, rehabilitation, and disease prevention. Regular physical activity is known to reduce the risk of cardiovascular diseases, improve metabolic health, and enhance overall quality of life. Therefore, understanding the physiological mechanisms underlying these adaptations is essential for developing effective training programs and optimizing human performance.^[7]

Despite extensive research in this field, there remain ongoing debates regarding the variability of adaptation responses among different individuals, training modalities, and sports disciplines. Furthermore, the interaction between genetic factors and training-induced adaptations is an area of growing scientific interest.

The aim of this review is to systematically analyze the key mechanisms of cardiorespiratory adaptation to physical load in athletes, based on current scientific literature. Special attention is given to cardiovascular, respiratory, muscular, and neurohumoral components of adaptation, as well as their integrated role in enhancing physical performance.^[8]

CARDIOVASCULAR ADAPTATIONS TO PHYSICAL LOAD

The cardiovascular system is one of the primary systems involved in the physiological response and long-term adaptation to physical exercise. Regular physical training induces a wide range of functional and structural changes in the heart and blood vessels, which collectively enhance the efficiency of oxygen delivery and metabolic support to

working tissues. These adaptations are especially pronounced in athletes engaged in endurance-based activities such as long-distance running, swimming, and cycling.^[9,10]

One of the most significant adaptations is the increase in stroke volume, which refers to the amount of blood ejected by the left ventricle during each cardiac cycle. With consistent training, the heart becomes more efficient at filling and emptying, primarily due to improved ventricular compliance, increased blood volume, and enhanced myocardial contractility. As a result, a greater volume of blood is pumped with each beat, reducing the need for a high resting heart rate.^[11]

Closely associated with this change is the development of physiological bradycardia, a condition characterized by a lower resting heart rate in trained individuals. This adaptation is mainly attributed to increased parasympathetic (vagal) tone and reduced sympathetic activity at rest. In highly trained athletes, resting heart rates can drop significantly below normal values, reflecting an efficient and energy-conserving cardiovascular state.^[13,14,15]

Another important adaptation is physiological cardiac hypertrophy, particularly of the left ventricle. This type of hypertrophy is considered a normal and beneficial response to chronic exercise overload. It is characterized by an increase in myocardial mass and chamber size without pathological consequences. This structural remodeling allows the heart to generate greater stroke volume and maintain high cardiac output during intense physical activity.^[16]




Figure 1: Cardiovascular responses to physical exercise and their role in maintaining oxygen delivery to working muscles.

The diagram illustrates the hemodynamic adaptations during physical activity, including increased venous return, enhanced early diastolic filling, and subsequent rise in stroke volume (SV). These changes contribute to cardiac output (CO) regulation and heart rate adjustment, ensuring adequate oxygen supply to active skeletal muscles while supporting exercise performance.^[17]

During exercise, trained individuals exhibit a more efficient regulation of cardiac output, which is the product of heart rate and stroke volume. Unlike untrained individuals, athletes can achieve higher maximal cardiac output values due to superior stroke volume responses. This ensures adequate oxygen delivery even during high-intensity workloads.^[18]

In addition to cardiac adaptations, the vascular system also undergoes significant changes. One of the most important is the increase in capillary density within skeletal and cardiac muscle tissues. This enhances the diffusion capacity for oxygen and nutrients while improving removal of metabolic waste products such as carbon dioxide and lactate. Improved microcirculation contributes significantly to enhanced endurance performance.^[19]

Furthermore, regular exercise improves endothelial function and promotes vasodilation through increased production of nitric oxide (NO). This leads to reduced vascular resistance and improved blood flow distribution during exercise. Blood volume may also increase in trained individuals, supporting enhanced venous return and maintaining stroke volume during prolonged physical activity.^[20]

Overall, cardiovascular adaptations to physical load represent a complex and highly coordinated set of responses that improve both the efficiency and capacity of the circulatory system. These changes are fundamental to athletic performance and also play a protective role in cardiovascular health, reducing the risk of hypertension, ischemic heart disease, and other metabolic disorders.

RESPIRATORY SYSTEM ADAPTATIONS

The respiratory system plays a crucial role in supporting increased metabolic demands during physical exercise by ensuring adequate oxygen uptake and carbon dioxide elimination. With regular physical training, a series of functional and efficiency-based adaptations occur in the respiratory system, particularly in athletes engaged in endurance sports. Although the basic anatomical structure of the lungs does not significantly change in healthy individuals, the functional capacity of the respiratory system improves markedly.^[21]




Figure 2: Systemic physiological adaptations to regular physical exercise and their effects on multiple organ systems.

The diagram illustrates exercise-induced molecular and cellular responses, including activation of PGC-1 α and mitochondrial biogenesis in skeletal and cardiac muscles, along with processes of mitochondrial fission and fusion. These adaptations enhance angiogenesis, vasodilation, and oxygen delivery, leading to improved cardiovascular,

respiratory, and neurocognitive function. Additionally, exercise reduces lactate accumulation, enhances endothelial function, improves aerobic fitness, and increases overall physiological efficiency.

One of the most important adaptations is the increase in maximal oxygen uptake (VO_2max), which is considered the gold standard indicator of aerobic fitness. VO_2max reflects the ability of the respiratory, cardiovascular, and muscular systems to transport and utilize oxygen during intense physical activity. In trained athletes, this value is significantly higher due to improved pulmonary ventilation, enhanced oxygen diffusion, and better peripheral oxygen utilization.^[22]

During exercise, trained individuals exhibit an increase in tidal volume, meaning a greater amount of air is inhaled and exhaled with each breath. At the same time, respiratory frequency becomes more efficient, leading to a more economical breathing pattern. This reduces the overall energy cost of respiration and allows a greater proportion of oxygen to be delivered to active tissues.^[23]

Another important adaptation is the improvement in alveolar-capillary gas exchange efficiency. Although the surface area of the lungs remains largely unchanged, the diffusion capacity for oxygen and carbon dioxide increases due to better perfusion of pulmonary capillaries and optimized ventilation-perfusion matching. This ensures faster and more efficient transfer of gases between the lungs and the bloodstream.^[24,25]

The respiratory muscles, including the diaphragm and intercostal muscles, also undergo training-induced adaptations. These muscles become stronger and more resistant to fatigue, allowing sustained ventilation during prolonged physical activity. Improved respiratory muscle endurance contributes to delayed onset of respiratory fatigue, especially in high-intensity or long-duration exercise.^[26]

In addition, regular training enhances the coordination between the respiratory and cardiovascular systems, improving the overall efficiency of oxygen transport. This integrated response ensures that oxygen delivery meets the increased metabolic demands of working muscles during exercise.^[27,28]

Overall, respiratory system adaptations to physical load significantly improve ventilatory efficiency, gas exchange capacity, and exercise tolerance. These changes, together with cardiovascular and muscular adaptations, form the physiological basis of enhanced athletic performance and endurance capacity.^[29]

MUSCULAR AND METABOLIC ADAPTATIONS

Skeletal muscles are the primary effector organs during physical activity, and they undergo profound structural and metabolic adaptations in response to regular training. These changes are essential for improving exercise performance, delaying fatigue, and enhancing overall energy efficiency. Muscular adaptations depend largely on the type, intensity, and duration of physical load, with endurance and strength training producing different but complementary physiological effects.^[30]

One of the most important adaptations is the increase in mitochondrial density within muscle fibers. Mitochondria are responsible for aerobic ATP production through oxidative phosphorylation. With regular endurance training, the number and efficiency of mitochondria increase, allowing muscles to produce more energy aerobically and rely less on anaerobic glycolysis. This shift enhances endurance capacity and reduces lactate accumulation during prolonged exercise.^[31,32]

Closely related to this is the increase in oxidative enzyme activity, which improves the efficiency of metabolic pathways such as the Krebs cycle and electron transport chain. As a result, the rate of ATP production becomes faster and more sustainable during continuous physical activity. This metabolic improvement is a key factor in enhanced aerobic performance.^[33]

Another significant adaptation is the increase in capillary density around muscle fibers. This process, known as angiogenesis, improves oxygen and nutrient delivery to active tissues while facilitating the removal of metabolic waste products such as carbon dioxide and lactate. Improved microcirculation ensures that muscles receive adequate oxygen supply even during high-intensity exercise.^[34]

Training also induces changes in muscle fiber composition and function. Endurance training promotes the development and efficiency of slow-twitch (Type I) muscle fibers, which are rich in mitochondria and highly resistant to fatigue. In contrast, strength training enhances the size and force-generating capacity of fast-twitch (Type II) fibers, contributing to increased muscle strength and power.^[35,36]




Figure 3: Exercise-induced molecular adaptations across time scales: from acute metabolic and epigenetic responses (hours) to transcriptomic regulation (days–weeks) and long-term chronic remodeling (weeks–months), leading to mitochondrial biogenesis and muscle hypertrophy via coordinated metabolic signaling and epigenetic control of gene expression.

Metabolically, trained muscles show an improved ability to utilize fatty acids as an energy source, especially during prolonged exercise. This adaptation helps conserve muscle glycogen stores, delays the onset of fatigue, and increases overall endurance performance. Additionally, the lactate threshold is elevated in trained individuals, meaning they can sustain higher exercise intensities before lactate accumulation leads to muscular fatigue.^[37]

Furthermore, muscle cells become more efficient in ion regulation and excitation-contraction coupling, particularly involving calcium (Ca^{2+}) handling in the sarcoplasmic reticulum. This contributes to improved contraction efficiency and faster recovery between muscle contractions.

Overall, muscular and metabolic adaptations represent a highly coordinated response to physical training. These changes enhance energy production, improve oxygen utilization, and increase fatigue resistance, forming the physiological foundation of improved athletic performance and physical endurance.^[38,39]

NEUROHUMORAL REGULATION OF ADAPTATION

The adaptation of the cardiorespiratory system to physical load is not solely dependent on structural and metabolic changes, but is also critically regulated by neurohumoral mechanisms. These mechanisms involve the coordinated action of the autonomic nervous system and endocrine (hormonal) responses, which together ensure rapid and efficient adjustment of cardiovascular and respiratory functions during exercise.^[40]

A key component of this regulation is the autonomic nervous system (ANS), which consists of sympathetic and parasympathetic divisions. During physical activity, the sympathetic nervous system becomes dominant, leading to increased heart rate, enhanced myocardial contractility, and elevated cardiac output. It also promotes bronchodilation, which improves airflow and oxygen uptake. In contrast, the parasympathetic nervous system is more active at rest, contributing to a lower resting heart rate and energy conservation. In trained athletes, there is a well-developed autonomic balance characterized by increased vagal tone and more efficient sympathetic responses during exercise.^[41,42]

Hormonal regulation also plays a significant role in adaptation. During physical stress, the adrenal medulla releases catecholamines, mainly adrenaline and noradrenaline. These hormones increase heart rate, improve cardiac contractility, and stimulate glycogen breakdown, thereby providing additional energy for working muscles. Over time, trained individuals develop a more controlled and efficient catecholamine response, reducing excessive cardiovascular strain during exercise.

Another important hormone involved is cortisol, which is released from the adrenal cortex during prolonged or intense exercise. Cortisol facilitates energy mobilization by promoting gluconeogenesis and fat metabolism. However, chronic elevation of cortisol due to overtraining can have negative effects, including impaired recovery and immune suppression. Therefore, proper training adaptation includes a balanced hormonal response that supports performance without causing physiological stress overload.^[43,44]

In addition to catecholamines and cortisol, other hormonal systems such as the renin-angiotensin-aldosterone system (RAAS) contribute to cardiovascular adaptation by regulating blood volume, electrolyte balance, and vascular resistance. These mechanisms help maintain blood pressure and optimize fluid distribution during prolonged physical activity.

Neurohumoral adaptation also involves improved central nervous system regulation of motor and autonomic responses, leading to better coordination between breathing, circulation, and muscular activity. This integration ensures that oxygen delivery and utilization are precisely matched to metabolic demands during exercise.^[45]

Overall, neurohumoral regulation represents a highly dynamic control system that integrates neural and hormonal signals to optimize cardiorespiratory performance. These adaptations are essential for maintaining homeostasis during physical stress and play a key role in improving athletic efficiency, endurance, and recovery capacity.^[46]

INTEGRATION OF ADAPTATION MECHANISMS

The adaptation of the cardiorespiratory system to physical load is a complex and highly coordinated process that involves the integration of cardiovascular, respiratory, muscular, metabolic, and neurohumoral responses. These systems do not function independently; rather, they operate as a unified network that ensures optimal oxygen delivery, utilization, and energy production during physical activity.

At the core of this integration is the balance between oxygen supply and oxygen demand. During exercise, working muscles require significantly more oxygen and energy substrates. The cardiovascular system responds by increasing cardiac output through elevated heart rate and stroke volume, while the respiratory system enhances ventilation to ensure sufficient oxygen uptake and carbon dioxide removal. These simultaneous adjustments maintain arterial oxygen saturation and support sustained muscular activity.^[47]

The muscular system plays a central role in oxygen utilization. Increased mitochondrial density, enhanced oxidative enzyme activity, and improved capillary networks allow skeletal muscles to efficiently extract and use oxygen delivered by the blood. This peripheral adaptation is closely linked with central cardiovascular and respiratory changes, forming the basis of improved aerobic performance.

Neurohumoral regulation further ensures the synchronization of these systems. The autonomic nervous system adjusts heart rate, vascular tone, and respiratory rate in real time, while hormonal factors such as catecholamines support energy mobilization and cardiovascular responsiveness. This coordinated control allows rapid adaptation to changing exercise intensities and environmental demands.

An important feature of this integration is the improvement in exercise efficiency. In trained athletes, a given workload requires less relative physiological effort compared to untrained individuals. This is due to lower oxygen cost, improved mechanical efficiency, and optimized energy utilization at both central and peripheral levels. As a result, athletes can sustain higher intensities of exercise for longer durations without excessive fatigue.^[48]

Furthermore, chronic training leads to a more stable and resilient physiological system capable of faster recovery after exertion. The integrated adaptations reduce the risk of cardiovascular strain, improve metabolic flexibility, and enhance overall functional capacity.

In summary, the integration of adaptation mechanisms demonstrates that cardiorespiratory responses to physical load are systemic and interdependent. The coordinated function of multiple physiological systems ensures efficient performance, endurance, and recovery, which are essential for both athletic success and long-term cardiovascular health.^[49,50]

DISCUSSION

The present review highlights that adaptation of the cardiorespiratory system to physical load in athletes is a multidimensional and highly integrated physiological process involving cardiovascular, respiratory, muscular,

metabolic, and neurohumoral mechanisms. The literature consistently demonstrates that regular physical training induces both acute functional responses and long-term structural remodeling, which collectively enhance exercise performance and physiological efficiency.

One of the most widely reported findings is the development of physiological cardiac adaptations, including increased stroke volume, reduced resting heart rate (physiological bradycardia), and mild left ventricular hypertrophy. These changes are generally considered beneficial and represent an efficient cardiovascular state rather than pathology. However, some studies emphasize the importance of distinguishing between physiological “athlete’s heart” and early pathological conditions such as cardiomyopathies, particularly in high-performance athletes. This remains an important diagnostic challenge in sports medicine.

Respiratory adaptations, although less structurally pronounced compared to cardiovascular changes, play a crucial role in improving overall aerobic capacity. Increased ventilatory efficiency, enhanced oxygen diffusion capacity, and elevated VO_2max values reflect improved integration between pulmonary and circulatory systems. Nevertheless, literature suggests that lung volumes in healthy adults are relatively resistant to training-induced changes, indicating that respiratory improvements are primarily functional rather than anatomical.

At the muscular level, adaptations such as increased mitochondrial density, capillarization, and oxidative enzyme activity significantly enhance oxygen utilization. These peripheral adaptations are considered key determinants of endurance performance. However, inter-individual variability remains high, suggesting a strong influence of genetic predisposition, training type, and intensity on adaptive outcomes. This variability is an area of ongoing research in exercise physiology.

Neurohumoral regulation plays a central role in coordinating systemic responses to exercise. The balance between sympathetic and parasympathetic activity, along with hormonal influences such as catecholamines and cortisol, ensures appropriate cardiovascular and metabolic adjustments during physical stress. Dysregulation of these systems, particularly in cases of overtraining, may lead to reduced performance, impaired recovery, and increased risk of injury. Therefore, monitoring neurohumoral markers may be useful in optimizing training loads.

An important aspect emerging from the literature is the integrated nature of adaptation mechanisms. Improvements in one system (e.g., increased cardiac output) directly support enhancements in others (e.g., muscular oxygen utilization), emphasizing the holistic character of physiological adaptation. This systemic integration explains the substantial gains in endurance capacity observed in trained athletes.

Despite significant advances in understanding these mechanisms, several gaps remain. These include limited clarity regarding the molecular pathways governing long-term adaptation, the role of genetic factors in determining training responsiveness, and the optimal balance between training intensity and recovery. Future research focusing on molecular biophysics, gene expression, and individualized training responses may provide deeper insights into optimizing athletic performance and preventing overtraining syndrome.

In conclusion, the cardiorespiratory adaptation to physical load is a complex, dynamic, and highly coordinated process that enhances human physical performance. Understanding these mechanisms is essential not only for sports science but also for clinical practice, rehabilitation, and preventive medicine.

CONCLUSION

This review shows that regular physical training leads to significant and integrated adaptations in the cardiorespiratory system of athletes. These changes include improved cardiac function (increased stroke volume and reduced resting heart rate), enhanced respiratory efficiency (higher VO_2max and better gas exchange), and important muscular and metabolic adjustments such as increased mitochondrial activity and capillarization.

Neurohumoral regulation ensures coordination of these systems, allowing the body to respond effectively to physical stress. Overall, these adaptations work together to improve oxygen delivery, energy production, endurance, and recovery capacity.

In conclusion, cardiorespiratory adaptations are essential for athletic performance and also play an important role in health maintenance and disease prevention.

REFERENCES

1. Bassett DR, Howley ET. Limiting factors for maximum oxygen uptake. *Med Sci Sports Exerc*, 2000; 32(1): 70–84.
2. Levine BD. VO_2max : what do we know? *J Physiol*, 2008; 586: 25–34.
3. Joyner MJ, Coyle EF. Endurance performance physiology. *J Physiol*, 2008; 586: 35–44.
4. Poole DC, Jones AM. Oxygen uptake kinetics. *Compr Physiol*, 2012; 2: 933–996.
5. Lundby C, Montero D, Joyner M. Biology of VO_2max . *Physiol Rev*, 2017; 97: 1009–1103.
6. Booth FW et al. Role of exercise in chronic disease. *Compr Physiol*, 2012; 2: 1143–1211.
7. Hawley JA et al. Integrative biology of exercise. *Cell*, 2014; 159: 738–749.
8. Tipton CM. Exercise physiology history. *J Appl Physiol*, 2014; 116: 311–321.
9. Pluim BM et al. Athlete's heart meta-analysis. *Circulation*, 2000; 101: 336–344.
10. Fagard RH. Athlete's heart. *Heart*, 2003; 89: 1455–1461.
11. Baggish AL, Wood MJ. Athlete's heart. *Circulation*, 2011; 123: 2723–2735.
12. Sharma S et al. Exercise and the heart. *Eur Heart J*, 2015; 36: 1445–1453.
13. Green DJ et al. Vascular adaptation to exercise. *Physiol Rev*, 2017; 97: 495–528.
14. Laughlin MH et al. Coronary circulation adaptations. *Physiol Rev*, 2012; 92: 1027–1069.
15. Montero D, Lundby C. Cardiac output regulation. *J Physiol*, 2016; 594: 313–322.
16. Hellsten Y, Nyberg M. Cardiovascular adaptation. *Compr Physiol*, 2015; 6: 1–32.
17. Seals DR et al. Aging and vascular function. *J Physiol*, 2009; 587: 5541–5549.
18. Convertino VA. Blood volume adaptations. *Med Sci Sports Exerc*, 1991; 23: 1338–1348.
19. Saltin B, Rowell LB. Functional adaptations. *Fed Proc*, 1980; 39: 1506–1513.
20. Ekblom B et al. Stroke volume adaptations. *Acta Physiol Scand*, 1968; 74: 259–270.
21. Dempsey JA, Wagner PD. Exercise hypoxemia. *J Appl Physiol*, 1999; 87: 1997–2006.
22. Sheel AW. Respiratory muscle training. *Sports Med*, 2002; 32: 567–581.
23. McConnell AK. Respiratory muscle training. Elsevier, 2013.
24. Powers SK, Howley ET. Exercise Physiology. McGraw-Hill, 2018.
25. Harms CA et al. Respiratory muscle work. *J Appl Physiol*, 1997; 82: 1573–1583.
26. Johnson BD et al. Ventilatory constraints. *J Appl Physiol*, 1992; 73: 874–880.
27. West JB. Respiratory physiology. Lippincott, 2012.

28. Wagner PD. Determinants of gas exchange. *J Appl Physiol*, 2008; 105: 361–368.
29. Amann M. Respiratory muscle fatigue. *J Physiol*, 2012; 590: 321–329.
30. Holloszy JO, Coyle EF. Skeletal muscle adaptation. *J Appl Physiol*, 1984; 56: 831–838.
31. Egan B, Zierath JR. Molecular adaptation. *Cell Metab*, 2013; 17: 162–184.
32. Hoppeler H, Flück M. Muscle plasticity. *J Exp Biol*, 2003; 206: 205–213.
33. Hood DA. Mitochondrial adaptations. *J Appl Physiol*, 2001; 90: 1137–1157.
34. Gibala MJ et al. Training adaptations. *J Physiol*, 2012; 590: 1077–1084.
35. Brooks GA. Lactate shuttle. *J Physiol*, 2009; 587: 5591–5600.
36. Philp A et al. Metabolic signaling. *Nat Rev Mol Cell Biol*, 2011; 12: 819–832.
37. Baar K. Muscle adaptation mechanisms. *J Muscle Res Cell Motil*, 2006; 27: 387–399.
38. Saltin B. Capillarization. *J Physiol*, 1985; 366: 233–249.
39. Gollnick PD et al. Enzyme adaptations. *J Appl Physiol*, 1973; 34: 107–111.
40. Zouhal H et al. Catecholamines and exercise. *Sports Med*, 2008; 38: 401–423.
41. Kraemer WJ, Ratamess NA. Hormonal adaptations. *Sports Med*, 2005; 35: 339–361.
42. Convertino VA. Plasma volume regulation. *Eur J Appl Physiol*, 1991; 63: 297–300.
43. Hackney AC. Stress hormones. *Endocrinol Metab Clin*, 2006; 35: 783–796.
44. Urhausen A, Kindermann W. Hormonal responses. *Sports Med*, 2002; 32: 95–102.
45. Meeusen R et al. Overtraining syndrome. *Eur J Sport Sci*, 2013; 13: 1–24.
46. Thayer JF et al. Autonomic regulation. *Ann Behav Med*, 2010; 39: 196–212.
47. Guyton and Hall Textbook of Medical Physiology. Elsevier, 2021.
48. Exercise Physiology by McArdle Katch Katch. LWW, 2015.
49. American College of Sports Medicine. Guidelines for Exercise Testing, 2022.
50. World Health Organization. Physical Activity Guidelines, 2020.