

HASHIMOTO'S ENCEPHALOPATHY PRESENTING WITH NEUROCOGNITIVE SYMPTOMS: A CASE REPORT

Zohair Ahmed¹ and Mohammed Maroof Mudassir*²

¹Final Year Part-II.

²Final Year Part-I, OSMEC-1585.

Article Received: 24 January 2025 | Article Revised: 13 February 2025 | Article Accepted: 06 March 2025

*Corresponding Author: Mohammed Maroof Mudassir

Final Year Part-I, OSMEC-1585.

DOI: <https://doi.org/10.5281/zenodo.15113844>

How to cite this Article: Zohair Ahmed and Mohammed Maroof Mudassir (2025). HASHIMOTO'S ENCEPHALOPATHY PRESENTING WITH NEUROCOGNITIVE SYMPTOMS: A CASE REPORT. World Journal of Pharmaceutical Science and Research, 4(2), 395-405. <https://doi.org/10.5281/zenodo.15113844>



Copyright © 2025 Mohammed Maroof Mudassir | World Journal of Pharmaceutical Science and Research.

This work is licensed under creative Commons Attribution-NonCommercial 4.0 International license (CC BY-NC 4.0)

ABSTRACT

A 45-year-old female presented with a one-month history of cough with expectoration, followed by intermittent fever, chest pain, and progressive neurological symptoms, including confusion and drowsiness. Her condition worsened with sudden-onset abdominal pain, vomiting, and bladder incontinence, prompting hospitalization. Initial assessments revealed accelerated hypertension, altered sensorium, and elevated anti-thyroid antibodies. Imaging showed old infarcts and intraparenchymal hemorrhage. After ruling out alternative causes, she was diagnosed with Hashimoto's Encephalitis, a rare autoimmune disorder characterized by cognitive impairment and neurologic deficits. She responded to glucocorticoid therapy and supportive treatment. This case highlights the importance of early recognition and intervention in autoimmune encephalitis.

KEYWORD: Intraparenchymal, cough, bladder, Hashimoto's Encephalitis.

CHIEF COMPLAINTS

Patient was brought by husband to the hospital with chief complaints of:

- Cough with expectoration since 30 days
- Fever since 5-6 days
- Chest pain since 5 days
- Confusion and drowsiness since 5-6 days

**PATIENT DETAILS**

Name- Mrs. Farhath Kauser Age- 45 Years

Sex- Female Informant- Self

**HISTORY OF PRESENTING
ILLNESS**

Patient was apparently asymptomatic 30 days ago. Patient had travelled to Saudi Arabia on pilgrimage following which she developed

- Cough with expectoration
- Chest pain which was not associated with fever.

1 week later, cough increased especially at night time with

- Intermittent low grade fever.

Since 2 days, patient presented with

- Altered sensorium
- Persistent drowsiness
- Accelerated HTN

From 5:30pm on 4/7/2024, patient developed

- Sudden pain abdomen
- An episode of vomiting
- Bladder incontinence. Then she was taken to nearby hospital and treatment was started.

OWAISI GROUP OF HOSPITAL
PRINCESS ESRA HOSPITAL
DECCAN COLLEGE OF MEDICAL SCIENCE

DISCHARGE CARD

Patient Name Mr./ Mrs. Faheem Khan
 S/o., W/o., D/o., Faheem Khan Age 45 Sex F
 I.P. No. 1346719 Unit. Neurology
 Consultant Name : Dr. Sandhya Manoj / Dr. Suman Kumar
 D.O.A. 5/7/2024 D.O.D. 6/7/2024

Diagnosis Acute Hemorrhagic Stroke
s/o. Acute Hemiparesis
acute bleed in the left Thalamocapsular

Chief Complaints Headache
BK- APSE LRTI
Keto on (uncontrolled) + Accelerated HTN

Present illness as per since 10 days.
cough since 10 days.

Past illness pt had travel history to Saudi Arabia
45 days ago. 5 days later she developed cough
& expectoration, not runny.
Swollen throat, cough increased at night time and had
intermittent low grade fever.

Family History 5 days, patient had increased drowsiness
from 5:30 pm on 4/7/24 - had sudden
pain abdomen, & 1 episode of vomiting and also

Drug History bladder incontinence. At hospital was
had incontinent then ⊕ and persistent drowsiness.

Investigation 180/110 mmHg
tb-40 hb-


PAST HISTORY

- History of Hypertension since 5 years
- History of Hypothyroidism since 8 years
- History Hair loss since one year
- No similar complaints in the past
- No H/O TB/Epilepsy/Asthma/COPD/ CAD/ Blood transfusions
- No H/O Head Trauma
- No H/O Major hospitalizations
- No H/O Major surgeries
- No H/O of other associated autoimmune disorders such as SLE/Vitiligo/Addison's disease.

FAMILY HISTORY

None of the patient's parents, siblings or first degree relatives have or have had similar complaints or any significant comorbidities.

CAUSES OF HYPOTHYROIDISM



Personal History

- Diet: Mixed
- Appetite: Normal
- Bowel & Bladder: Irregular
- Sleep: Disturbed
- Addictions: None
- Drug History: Patient was taking Regesterone

GENERAL EXAMINATION


- On examination, the 45-year-old patient was in altered sensorium with persistent drowsiness, is moderately built, well-nourished and conscious, cooperative, and comfortably lying on the bed, oriented to time, place and person.
- Puffy Face w/ edematous eyelids.
- Non-pitting pre-tibial edema (Myxedema) seen.
- Goitre: Firm and rubbery in consistency (Characteristic).
- There is No Pallor, Icterus, cyanosis, koilonychia and no generalized lymphadenopathy.

Vitals

- Temperature: 101 °F
- Blood Pressure: 180/110 mmHg
- Pulse Rate: Tachycardia 80 beats/min
- SpO₂: 96% on room air
- RR: 19-21 breaths/min

Patient Details

- Height: Approx 5'3
- Weight: 63kg



SYSTEMIC EXAMINATION

1. Intellectual Functions

- Handedness: Right
- Memory: Recent and Remote intact
- Appearance: Well-kept
- GCS Score- E2V5M6
- Speech: Normal

2. Cranial Nerves

<u>Nerve</u>	Procedure	Right and Left
<u>1. Olfactory</u>	Close eyes, close one nostril and repeat on both sides. Use smelling salt, coffee, asafetida	Normal
<u>2. Optic</u>	1. Visual acuity: Snellen/ Counting Fingers from 6m onwards, Hand Movements close to face, Projection of light and Perception of light 2. Visual Field: (Confrontation) 3. Colour vision 4. Reflexes- Light and Accom 5. Fundus Exam	Normal
<u>3,4,6 CN</u>	Ocular movements	Normal
<u>Trigeminal Nerve</u>	Motor: Pterygoids, Temporalis (Mastication) Sensory: Over face, corneal, conjunctival reflex, jaw jerk	Normal
<u>Facial Nerve</u>	Deviation of mouth, frowning, closure of eyes against resistance, blowing, whistling, smiling, accumulation of tears, drooling of saliva	B/L Abductor paresis - facial asymmetry
<u>Vestibulocochlear Nerve</u>	Whispering, watch test, rinnes, weber, schwabach	Normal
<u>Vagus & Glossopharyngeal</u>	Gag reflex, uvula position	Normal
<u>Spinal Accessory</u>	Shrug shoulders, movement of head to one side against resistance	Normal
<u>Hypoglossus</u>	Deviation of tongue, See tongue inside the mouth for fasciculations	Normal

3. Motor System

A. Attitude and Position: Normal

B. Nutrition (Bulk) - Normal

C. Increased tone:

Neck Stiffness +

Rt pronator drift +

D. Power: UL LL, Right Left

4/5, 4/5

E. Abnormal Movements: -

4. REFLEXES: [For both sides]

Superficial: 2+ (Normal)

Deep: B/L Plantar Flexion +

5. SENSORY SYSTEM

Superficial: Normal

Deep: Normal

6. CEREBELLUM

Ataxia + (Rt, Lt)

7. COORDINATION AND GAIT

Ataxic Gait +

8. Skull and Spine: Normal**9. Signs of Meningeal Irritation: -****CASE SUMMARY**

Over a months period, a 45-year-old female who was previously healthy developed cough with expectoration which was initially not associated with fever. One week later, cough increased at night time and had intermittent low-grade fever. The patient reported slowness in performing her daily activities, walking difficulties and trouble with getting changed. Additionally, her relatives observed transient episodes of fluctuating consciousness with drowsiness.

Initially the patient was admitted to Princess Esra Hospital where she was found to have increased drowsiness. On 4/7/24 from 5:50pm onwards patient suddenly developed pain abdomen with 1 episode of vomiting and bladder incontinence. BP at the hospital was accelerated above 180mmHg systolic. Patient was administered anti-seizure medication, anti-hypertensives and supportive treatment.

Laboratory studies at the time did not yield any significant findings except for mild leucocytosis. After a few days, patient's attendants refused further management and willing to go to a different hospital. Patient was advised Gynaecology opinion as patient was taking Regesterone at the time.

Patient underwent multiple investigations including LFT, RFT, CT Brain (Plain), etc at the latter hospital and was found to have positive findings. For example, her Anti TPO was excessively elevated following which she was diagnosed as a case of Hashimoto's Encephalitis after ruling out other diseases.

Provisional Diagnosis: Suspected Encephalitis following URTI with Hypothyroidism and Accelerated HTN.

PATHOPHYSIOLOGY

Hashimoto's Encephalitis (HE) is an unusual neurologic disorder whose etiology, pathogenesis and histologic characteristics are unclear.

A hospital-based epidemiologic study of neurologic symptoms consistent with HE estimated its prevalence to be about 2.1 per 100,000. The disorder occurs more frequently between age 44 to 46 years, with a female to-male ratio of four to one.

The clinical manifestations usually include acute to subacute onset of confusion with alteration of consciousness. Two major patterns of presentation were described: (1) 25% of patients follow a stroke-like pattern of multiple recurrent

episodes of focal neurologic deficits with a variable degree of cognitive dysfunction and consciousness impairment, (2) and the remaining 75% present with a diffuse progressive pattern of slow cognitive decline with dementia, confusion and hallucinations. These two clinical patterns may overlap over the course of the disease. In this case report, our patient's clinical manifestations are more consistent with the first form of presentation, which is less common.

The pathogenic role of thyroid antibodies remains unknown, there is no evidence that any antithyroid anti-body reacts with brain tissue or affects nerve function, and there is no clear correlation between the severity of the neurologic symptoms and the concentration of these antibodies.

Hashimoto's encephalopathy has been generalised as a steroid-responsive syndrome associated with TPO antibodies, myoclonus, and a slow-wave activity on electroencephalography.

DIFFERENTIAL DIAGNOSIS

The differential diagnosis of HE must consider any condition associated with delirium, rapidly progressive dementia, seizures or focal neurologic deficits. Thus, the list of diseases that can be confused with HE is vast, including stroke or transient ischemic attack, cerebral vasculitis, carcinomatous meningitis, toxic metabolic encephalopathies, paraneoplastic syndromes, Creutzfeldt-Jakob disease, degenerative dementia and psychiatric diseases.



PROGNOSIS

The long-term prognosis is variable, although a high percentage of patients respond to treatment; others could have a progressive or a relapsing course. The symptoms usually improve with glucocorticoid therapy however, it is not necessary because of treatment. A systematic review of 85 cases published on HE found clinical response in 98% of patients treated with glucocorticoids, 92% of patients treated with glucocorticoids and levothyroxine and 67% of patients treated with levothyroxine only.

1) CBP

RBC count: 4.9 million/cumm

WBC count: 12100 cells/cumm

Platelets: 4.3 Lakhs/cumm

Haemoglobin: 10.1 gms%

Differential Count

- Neutrophils: 75%
- Lymphocytes: 15%
- Monocytes: 06%
- Eosinophils: 04%
- Basophils: 00%

Impression

WBC: Leucocytosis,

RBC: Normocytic Normochromic Anisocytosis,

PLT: Adequate.

2) RFT

Urea: 27 mg/dl

Creatinine: 0.9 mg/dl

Uric acid: 4.5 mg/dl

INVESTIGATIONS

3) LFT: Serum Albumin: 3.9g/dl

AST: 13 U/L

ALT: 16 U/L

4) Serum Electrolytes

Sodium: 135.4 mmol/L

Potassium: 3.89 mmol/L

Chloride: 98.2 mmol/L

5) Immunoassay

- Anti Thyroglobulin Antibody (Anti-TG): 695.34 IU/ml (normal: 0-4.11 IU/ml)
- Anti Thyroperoxidase Antibodies (Anti-TPO): 214.3


IU/ml (Normal: less than 50)

6) Glycosylated Haemoglobin A1C (HbA1C): 8.0%

Non-Diabetic <=5.8%

Pre-Diabetic 5.8-6.5%

Diabetic >= 6.5%

7) CT BRAIN (PLAIN)**Impression**

- Old infarcts are seen in the bilateral centrum semiovale and periventricular regions.
- Intraparenchymal hemorrhage in the left ganglio-thalamic region.

FINAL DIAGNOSIS

Hashimoto's
Encephalopathy

MANAGEMENT

- Tab. Prednisolone (20mg) - 2 days TID
10 days BD
20 days OD
- Tab. Quetiapine (25mg) - for sleep PRN
- Tab. Cilnidipine (5mg) - BD

- Tab. Ursodeoxycholic Acid (300mg) - BD
- Tab. Telmisartan (40mg) + Metoprolol Succinate (50mg) - BD
- Tab. Thyroxine (50mcg) - OD
- Tab. Levetiracetam (500mg) - BD
- Inj. Insulin Lispro (100IU/ml) – TID
- Tab. Rosuvastatin (5mg) - OD x 30days
- Tab. Citicoline (500mg) - BD x 60days

Supportive Treatment

- Inj. Methylcobalamin (500mcg)
- Tab. Calcium (200mg) + Vitamin D3 (0.25mcg) - OD x 30days
- Flexabenz Gel with Diclofenac Diethylamine & Menthol - PRN

UNDER THE GUIDANCE OF

Department of General Medicine (Deccan College of Medical Sciences, Owaisi Group of Hospitals):

Dr. Abubaker, MD Gen Med (HOD)

REFERENCES

1. Harrison's Principles of Internal Medicine, 21st edition.
2. Chong J, Rowland L, Utiger R: Hashimoto encephalopathy: syndrome or myth? Arch Neurol, 2003; 60: 164-71.
3. Peschen-Rosin R, Schabet M, Dichgans J: Manifestation of Hashimoto's encephalopathy years before onset of thyroid disease. Eur Neurol, 1999; 41: 79-84.
4. Canelo-Aybar et al.: Hashimoto's encephalopathy presenting with neurocognitive symptoms: a case report. Journal of Medical Case Reports, 2010; 4: 337.